

Treewidth of grid subsets

Eli Berger^{*} Zdeněk Dvořák[†] Sergey Norin[‡]

Abstract

Let Q_n be the graph of $n \times n \times n$ cube with all non-decreasing diagonals (including the facial ones) in its constituent $1 \times 1 \times 1$ cubes. Suppose that a set $S \subseteq V(Q_n)$ separates the left side of the cube from the right side. We show that S induces a subgraph of tree-width at least $\frac{n}{\sqrt{18}} - 1$. We use a generalization of this claim to prove that the vertex set of Q_n cannot be partitioned to two parts, each of them inducing a subgraph of bounded tree-width.

Let G_n be the plane triangulated $n \times n$ grid, and consider any (non-proper) coloring of vertices of G_n by two colors. A well-known HEX lemma implies that G_n contains a monochromatic path with at least n vertices. We consider a 3-dimensional analogue of this claim.

Let Q_n be the $n \times n \times n$ grid with all non-decreasing diagonals in its constituent unit cubes; more precisely, the vertex set of Q_n is $\{(x, y, z) \in \mathbb{Z}^3 : 0 \leq x, y, z \leq n - 1\}$ and two distinct vertices (x, y, z) and (x', y', z') are adjacent if $x \leq x' \leq x + 1$, $y \leq y' \leq y + 1$ and $z \leq z' \leq z + 1$. A result of Matoušek and Průvřetivý [12] applied in this special case shows that any coloring of vertices of Q_n by two colors contains a connected monochromatic subgraph with $\Omega(n^2)$ vertices. We aim to show that it actually contains a large monochromatic subgraph which is “2-dimensional” in nature, i.e., with a large grid minor. It is well-known that a graph contains a large grid as a minor if and only if it has a large tree-width [3, 4, 14, 15], and thus we can state our main result in the following equivalent form.

^{*}University of Haifa, Haifa, Israel. Supported by BSF grant no. 2006099 and by ISF grant no. 1581/12.

[†]Charles University, Prague, Czech Republic. E-mail: rakdver@iuuk.mff.cuni.cz. Supported by the Center of Excellence – Inst. for Theor. Comp. Sci., Prague, project P202/12/G061 of Czech Science Foundation and by the project LL1201 (Complex Structures: Regularities in Combinatorics and Discrete Mathematics) of the Ministry of Education of Czech Republic.

[‡]Department of Mathematics and Statistics, McGill University. Email: snorin@math.mcgill.ca. Supported by an NSERC grant 418520.

Theorem 1. *For every $t \geq 0$, there exists $n \geq 1$ such that for any partition A_1, A_2 of the vertex set of Q_n , either A_1 or A_2 induces a subgraph of Q_n of tree-width at least t .*

Recall that a *tree decomposition* (T, β) of a graph G is a tree T and a function $\beta : V(T) \rightarrow 2^{V(G)}$ assigning a *bag* $\beta(u)$ to each vertex $u \in V(T)$, such that every vertex of G , as well as both ends of every edge of G , are contained in some bag, and such that $\{u : v \in \beta(u)\}$ induces a connected subtree of T for every $v \in V(G)$. The *width* of the decomposition is the maximum of the sizes of its bags minus one, and the *tree-width* $\text{tw}(G)$ of G is the minimum of the widths of its tree decompositions.

Let us remark that the presence of diagonals is important for the validity of Theorem 1; if the diagonals of Q_n are omitted, the graph becomes bipartite, and thus it can be partitioned to two independent sets (of tree-width 0).

Theorem 1 is motivated by a notion from the algorithmic graph theory. We say that a class $\mathcal{G}$ of graphs is *tree-width fragile* if for every $k \geq 1$, there exists $t_k \geq 0$ such that every graph $G \in \mathcal{G}$, there exist pairwise disjoint sets $A_1, \dots, A_k \subseteq V(G)$ satisfying $\text{tw}(G - A_i) \leq t_k$ for $1 \leq i \leq k$. For example, planar graphs are known to have this property [2, 13]. Many interesting graph problems have efficient algorithms when restricted to graphs with bounded tree-width, enabling Baker [2] to exploit this property in design of approximation algorithms for planar graphs.

It is natural to ask which more general graph classes are tree-width fragile, as the same approximation algorithms can be used for such graph classes. Eppstein [10] proved that this is the case for graphs avoiding some apex graph as a minor (a graph H is *apex* if $H - v$ is planar for some $v \in V(H)$), and in particular for graphs embedded in any fixed surface. DeVos et al. [6] generalized the argument to all proper minor-closed classes of graphs.

Of course, not all graph classes are tree-width fragile. It is easy to see that any graph class with this property must be sparse and must have sublinear separators [8]. Theorem 1 gives another class of obstructions.

Corollary 2. *The graph class $\{Q_n : n \geq 1\}$ is not tree-width fragile.*

Indeed, Theorem 1 shows that the condition of tree-width fragility fails already for $k = 2$; unlike the previously known graph classes that are not tree-width fragile, the graphs Q_n have bounded maximum degree and balanced separators of order $O(|V(Q_n)|^{2/3})$. Interestingly, the class $\{Q_n : n \geq$

1} is *fractionally* tree-width fragile, where the fractional version of tree-width fragility is defined in the standard way [8]; it is the first known class of graphs showing that tree-width fragility and fractional tree-width fragility do not coincide.

Another motivation for Theorem 1 comes from graph coloring theory. Many results are known on the variants of the coloring where the color classes are not required to be independent sets, but rather satisfy some other constraints, such as inducing subgraphs of bounded maximum degree [5, 9], with bounded component size [1, 11], or, as in our case, bounded tree-width [6, 7]. For this notion of *low tree-width coloring*, the previous results mostly focus on positive results, showing that graphs from some class have a low tree-width coloring using a constant number of colors. On the other hand, Theorem 1 gives a lower bound, presenting an example of a natural class of graphs that do not have low tree-width coloring by two colors.

Let us now give a brief idea of the proof of Theorem 1. Recall that every graph containing a $t \times t$ grid as a minor has tree-width at least t . Hence, we aim to construct a monochromatic grid in Q_n by connecting appropriately chosen connected subgraphs by disjoint paths. Of course, we need to deal with the situation that such paths are blocked by the vertices of the other color. Such blocking subgraphs must be in a sense larger than the subgraphs we are trying to join. We now switch to the other color class, considering these blocking subgraphs to be the nodes of a grid we are constructing and trying to find paths between these nodes. We may switch back and forth between the color classes several times, repeatedly enlarging the node subgraphs. For this procedure to end, we need to argue that eventually, the node subgraphs cannot be separated by a set inducing a subgraph of small tree-width. Abstracting the problem further, we reach the following claim of independent interest.

Let X be a subset of vertices of Q_n that separates the left side of the grid from the right side. Supposing that X is minimal with this property, a geometric intuition tells us that X should correspond to a surface in the 3-dimensional Euclidean space separating the left side of Q_n from the right side, and that such a surface should contain a subdivision of a large 2-dimensional grid in Q_n . Although this geometric intuition is somewhat misleading and difficult to make precise, the overall conclusion that $Q_n[X]$ should have large tree-width is true.

Theorem 3. *Let $n \geq 1$ be an integer, let S_1 be the set of vertices of the left side of Q_n and let S_2 be the set of vertices of the right side of Q_n . If a set*

$X \subseteq V(Q_n) \setminus (S_1 \cup S_2)$ intersects every paths in G from S_1 to S_2 , then the subgraph of Q_n induced by X has tree-width at least $\frac{n}{\sqrt{18}} - 1$.

Again, note that the presence of diagonals is necessary for the validity of Theorem 3. To prove Theorem 1, we need a generalization of Theorem 3. A plane graph H is a *near-triangulation* if every face of H except for the outer one has length three. A triple (G, S_1, S_2) , where G is a graph and S_1 and S_2 are vertex-disjoint connected subgraphs of G , is an $(n \times n)$ -*slab* (with sides S_1 and S_2) if there exist pairwise vertex-disjoint near-triangulations $R_1, \dots, R_n \subseteq G$ (the *rows* of the slab) and pairwise vertex-disjoint near-triangulations $C_1, \dots, C_n \subseteq G$ (the *columns* of the slab) such that for $1 \leq i, j \leq n$, the intersection $R_i \cap C_j$ is a path with one end in S_1 and the other end in S_2 , and such that for $s \in \{1, 2\}$ and for every row or column H , the intersection of H with S_s is a subpath of the boundary of the outer face of H .

Theorem 4. *Let (G, S_1, S_2) be an $(n \times n)$ -slab with rows and columns of maximum degree at most $\Delta \geq 3$ and let X be a subset of $V(G) \setminus (V(S_1) \cup V(S_2))$. If every path in G from S_1 to S_2 intersects X , then $G[X]$ has tree-width at least $\frac{n}{\sqrt{3\Delta}} - 1$.*

Clearly, Theorem 4 implies Theorem 3. The proof of Theorem 4 is topological in nature and we give it in the following two sections. Section 3 is devoted to the proof of Theorem 1.

1 $\{0, \pm 1, \star\}$ -valued functions on graphs and homotopy

In this section, we develop a discrete variant of the basic tools of homotopy theory.

Let $\mathcal{L} = \{0, \pm 1, \star\}$. We will consider function $f : V(G) \rightarrow \mathcal{L}$ for a graph G . Recall that a *separation* of G is pair (A, B) of subsets of vertices such that $V(G) = A \cup B$ and G has no edge with one end in $A \setminus B$ and the other end in $B \setminus A$. Without explicitly assuming it, we will typically consider the vertices with function values -1 and 1 as corresponding to sets $A \setminus B$ and $B \setminus A$, respectively, for some separation (A, B) of G , and label 0 , as roughly corresponding to a union of some components of $G[A \cap B]$. This motivates the following definition. We say that $f : V(G) \rightarrow \mathcal{L}$ is *continuous* if vertices $u, v \in V(G)$ are not adjacent whenever $f(v) = 1$ and $f(u) = -1$. We say that $f : V(G) \rightarrow \mathcal{L}$ is *holomorphic* if f is continuous and additionally

vertices $u, v \in V(G)$ are not adjacent whenever $f(v) = 0$ and $f(u) = \star$. We say that $f : V(G) \rightarrow \mathcal{L}$ is *entire*, if it is continuous and $\star \notin \text{Image}(f)$.

Let $C^0(G, \mathcal{L})$ denote the set of functions $f : V(G) \rightarrow \mathcal{L}$. Fix an arbitrary orientation of $E(G)$, that is for every $e \in E(G)$ we distinguish its beginning vertex denoted by e^- and its end vertex denoted by e^+ . Let $C^1(G, \mathbb{Z})$ denote the lattice of all functions $f : E(G) \rightarrow \mathbb{Z}$. The operator $d : C^0(G, \mathcal{L}) \rightarrow C^1(G, \mathbb{Z})$ is defined by $df(e) = f(e^+) - f(e^-)$ if $\star \notin \{f(e^+), f(e^-)\}$ and $df(e) = 0$, otherwise.

For a directed walk $W = (v_0, e_1, v_1, \dots, e_n, v_n)$ on G , not necessarily respecting the orientation we fixed, and for $1 \leq i \leq n$, let $\varepsilon_i \in C^1(G, \mathbb{Z})$ be defined by $\varepsilon_i(e_i) = 1$ if $v_i = e_i^+$, $\varepsilon_i(e_i) = -1$ if $v_i = e_i^-$ and $\varepsilon_i(e) = 0$ for $e \in E(G) \setminus \{e_i\}$. Let $I_W = \sum_{i=1}^n \varepsilon_i$. For a walk W and $h \in C^1(G, \mathbb{Z})$ we define $\int_W h := \sum_{e \in E(G)} I_W(e)h(e)$. Let W^{-1} denote the reversal of W , that is the walk $(v_n, e_n, v_{n-1}, e_{n-1}, \dots, e_1, v_0)$. For two walks W_1 from u to v and W_2 from v to w , let $W_1 W_2$ denote the concatenation of W_1 and W_2 .

Lemma 5. *If W is a walk from u to v on a graph G and $f \in C^0(G, \mathcal{L})$ is entire on $V(W)$ then*

$$\int_W df = f(v) - f(u).$$

Proof. Let $W = (v_0, e_1, v_1, \dots, e_n, v_n)$, where $v_0 = u$ and $v_n = v$, and for $1 \leq i \leq n$, let ε_i be as in the definition of I_W . We have

$$\begin{aligned} \int_W df &= \sum_{e \in E(G)} I_W(e)df(e) = \sum_{i=1}^n \varepsilon_i(e_i)df(e_i) \\ &= \sum_{i=1}^n f(v_i) - f(v_{i-1}) = f(v_n) - f(v_0) = f(v) - f(u). \end{aligned}$$

□

We refer to closed walks of length three on a graph G as *triangles*. We say that a triangle T is *f-contractible* for $f \in C^0(G, \mathcal{L})$, if f is holomorphic on $V(W)$. The next observation is the first step towards obtaining an extension of Lemma 5 to non-entire functions.

Lemma 6. *Let T be a triangle in G and let $f \in C^0(G, \mathcal{L})$. If f is continuous then $|\int_T df| \leq 1$, and if T is f -contractible then $\int_T df = 0$.*

Proof. Let $V(T) = \{u, v, w\}$. If f is entire on $V(T)$, then the claim follows from Lemma 5. Hence, we can assume that $f(w) = \star$. If $f(u) = \star$ or $f(v) = \star$, then $df(e) = 0$ for every $e \in E(T)$ and $\int_T df = 0$.

Otherwise, $|\int_T df| = |f(u) - f(v)|$. Since f is continuous, we have $|f(u) - f(v)| \leq 1$. Furthermore, if T is f -contractible, then $f(u) \neq 0 \neq f(v)$, and thus $f(u) = f(v)$ and $\int_T df = 0$. $\square$

Let $f \in C^0(G, \mathcal{L})$ be continuous. From Lemma 6 it follows that $\int_W df = 0$ for every W that is a sum of f -contractible triangles. We need a slight generalization of this fact. We will say that a closed directed walk W on G is (f, k) -almost contractible if there exist triangles $T_1, T_2, \dots, T_n$ of G such that $I_W = \sum_{i=1}^n I_{T_i}$ and $T_{k+1}, T_{k+2}, \dots, T_n$ are f -contractible.

Let W_i be a walk from u_i to v_i on G for $i = 1, 2$. We say that W_1 and W_2 are (f, k) -almost homotopic if there exists a walk Q from u_1 to u_2 and a walk R from v_1 to v_2 such that the walk $QW_2R^{-1}W_1^{-1}$ is (f, k) -almost contractible, and furthermore f is constant and integer on each of $V(Q)$ and $V(R)$. We say that W_1 and W_2 are f -homotopic if they are $(f, 0)$ -almost homotopic. From Lemma 6 we deduce the following corollary.

Lemma 7. *Let $f \in C^0(G, \mathcal{L})$ be continuous and let W_1 and W_2 be walks on G . If W_1 and W_2 are (f, k) -almost homotopic then*

$$\left| \int_{W_1} df - \int_{W_2} df \right| \leq k.$$

In particular, if W_1 and W_2 are f -homotopic then $\int_{W_1} df = \int_{W_2} df$.

Proof. Let Q and R be the walks and $T_1, T_2, \dots, T_n$ be the triangles showing that W_1 and W_2 are (f, k) -almost homotopic. Let $W = QW_2R^{-1}W_1^{-1}$. By Lemma 6, we have

$$\begin{aligned} \left| \int_W df \right| &= \left| \sum_{e \in E(G)} I_W(e) df(e) \right| = \left| \sum_{e \in E(G)} \sum_{i=1}^n I_{T_i}(e) df(e) \right| \\ &= \left| \sum_{i=1}^n \int_{T_i} df \right| = \left| \sum_{i=1}^k \int_{T_i} df \right| \leq k. \end{aligned}$$

On the other hand, since f is constant on Q and R , Lemma 5 implies that

$$\left| \int_W df \right| = \left| \int_Q df + \int_{W_1} df - \int_R df - \int_{W_2} df \right| = \left| \int_{W_1} df - \int_{W_2} df \right|,$$

and the claim of the lemma follows. $\square$

For a function $h : Y \rightarrow \mathbb{R}$ and a set $X \subseteq Y$, let $h(X) = \sum_{x \in X} h(x)$. Let $P = v_0 v_1 \dots v_n$ be a path in a graph G . For a function $f \in C^0(G, \mathcal{L})$ that is entire on P and for $1 \leq i \leq n-1$, let $\lambda_{P,f}(v_i) = \frac{1}{2}(f(v_{i+1}) - f(v_{i-1}))$.

Lemma 8. *Let $f \in C^0(G, \mathcal{L})$ be a function that is entire on a path $P = v_0 v_1 \dots v_n$ in a graph G , such that $f(v_0) \neq 0$ and $f(v_n) \neq 0$. Suppose that $g \in C^0(G, \mathcal{L})$ satisfies $g(v) = f(v)$ for every $v \in V(P)$ such that $g(v) \neq \star$, and $f(v) = 0$ for every $v \in V(P)$ such that $g(v) = \star$. Let $X = \{v \in V(P) : g(v) = 0\}$. Then*

$$\frac{1}{2} \int_P dg = \lambda_{P,f}(X).$$

Furthermore, if g is holomorphic on P , then $\lambda_{P,f}(X)$ is an integer.

Proof. Let $P_1, \dots, P_k$ be the maximal subpaths of P such that g is entire on P_i for $i = 1, \dots, k$. Note that

$$\int_P dg = \sum_{j=1}^k \int_{P_j} dg,$$

and thus it suffices to prove that $\frac{1}{2} \int_{P_j} dg = \lambda_{P,f}(X \cap V(P_j))$ for $1 \leq j \leq k$. Let $P_j = v_a v_{a+1} \dots v_b$. By Lemma 5, we have $\frac{1}{2} \int_{P_j} dg = \frac{1}{2}(g(v_b) - g(v_a)) = \frac{1}{2}(f(v_b) - f(v_a))$. Let $V(P_j) \cap X = \{v_{c_1}, v_{c_2}, \dots, v_{c_p}\}$, where $c_1 < c_2 < \dots < c_p$. Since f is continuous on P , for $1 \leq i \leq p-1$ we have $f(v_{c_i+1}) = f(v_{c_i+1-1})$. Furthermore, $f(v_{c_1-1}) = f(v_a)$ and $f(v_{c_p+1}) = f(v_b)$; this is the case even if say $c_1 = a$, since then $g(v_{a-1}) = \star$ and $f(v_{a-1}) = f(v_a) = 0$. Therefore, $\lambda_{P,f}(X \cap V(P_j)) = \frac{1}{2} \sum_{i=1}^p f(v_{c_i+1}) - f(v_{c_i-1}) = \frac{1}{2}(f(v_b) - f(v_a)) = \frac{1}{2} \int_{P_j} dg$, as required. Moreover, if g is holomorphic on P , then $f(v_a) = \pm 1$, since either $a = 0$ or $g(v_{a-1}) = \star$, and similarly $f(v_b) = \pm 1$. Hence, $\int_{P_j} dg \in \{-2, 0, 2\}$ and $\lambda_{P,f}(X \cap V(P_j)) = \frac{1}{2} \int_{P_j} dg$ is an integer. $\square$

2 Treewidth of slab separators

To give a lower bound on tree-width, we use the following claim (which is well-known, although usually stated with only non-negative weights).

Lemma 9. *Let $t \geq 0$ be an integer. Let H be a graph with $\text{tw}(H) \leq t$. Let $\lambda : V(H) \rightarrow \mathbb{R}$ be such that $|\lambda(v)| \leq 1$ for every $v \in V(H)$. If $\lambda(V(H)) \geq 3t + 3$, then there exists a separation (K, L) of H such that*

$$\frac{1}{3} \lambda(V(H)) \leq \lambda(K \setminus L) \leq \frac{2}{3} \lambda(V(H))$$

and $|K \cap L| \leq t + 1$.

Proof. Let (T, β) be a tree decomposition of H with bags of size at most $t+1$. For an edge uv of T , let $T_{u,v}$ denote the component of $T - uv$ containing v , and let $S_{u,v} = \bigcup_{w \in V(T_{u,v})} \beta(w) \setminus \beta(u)$.

We first show that there exists $u \in V(T)$ such that every neighbor v of u in T satisfies $\lambda(S_{u,v}) \leq \frac{2}{3}\lambda(V(H))$. If not, then for each $u \in V(T)$, let $\pi(u)$ denote a neighbor of u in T such that $\lambda(S_{u,\pi(u)}) > \frac{2}{3}\lambda(V(H))$. Since T is a tree, there exists an edge $uv \in E(T)$ such that $\pi(u) = v$ and $\pi(v) = u$. However, then

$$\lambda(V(H)) = \lambda(S_{u,v}) + \lambda(S_{v,u}) + \lambda(\beta(u) \cap \beta(v)) > \frac{4}{3}\lambda(V(H)) - (t+1),$$

which contradicts the assumption $\lambda(V(H)) \geq 3t+3$.

Let $u \in V(T)$ be a vertex such that every neighbor v of u in T satisfies $\lambda(S_{u,v}) \leq \frac{2}{3}\lambda(V(H))$. Let $v_1, v_2, \dots, v_m$ be the neighbors of u in T ordered so that $\lambda(S_{u,v_1}) \geq \lambda(S_{u,v_2}) \geq \dots \geq \lambda(S_{u,v_m})$. Note that $\sum_{i=1}^m \lambda(S_{u,v_i}) = \lambda(V(H)) - \lambda(\beta(u)) \geq \lambda(V(H)) - (t+1) \geq \frac{1}{3}\lambda(V(H))$. Let m' be the smallest index such that

$$\sum_{i=1}^{m'} \lambda(S_{u,v_i}) \geq \frac{1}{3}\lambda(V(H)).$$

Let $K = \beta(u) \cup \bigcup_{i=1}^{m'} S_{u,v_i}$ and $L = \beta(u) \cup \bigcup_{i=m'+1}^m S_{u,v_i}$. By the choice of m' , we have $\lambda(K \setminus L) \geq \frac{1}{3}\lambda(V(H))$. If $m' = 1$, then $\lambda(K \setminus L) = \lambda(S_{u,v_1}) \leq \frac{2}{3}\lambda(V(H))$. Hence, we can assume that $m' \geq 2$, and thus $\lambda(S_{u,v_i}) < \frac{1}{3}\lambda(V(H))$ for $1 \leq i \leq m$. By the choice of m' , we have $\bigcup_{i=1}^{m'-1} \lambda(S_{u,v_i}) < \frac{1}{3}\lambda(V(H))$, and thus $\lambda(K \setminus L) < \frac{1}{3}\lambda(V(H)) + \lambda(S_{u,v_{m'}}) < \frac{2}{3}\lambda(V(H))$. $\square$

We are now ready to prove our first main result.

Proof of Theorem 4. Let $R_1, \dots, R_n$ and $C_1, \dots, C_n$ be rows and columns of G of maximum degree at most Δ . For $1 \leq i, j \leq n$, let $G \uparrow (i, j)$ denote the path $R_i \cap C_j$, directed from S_1 to S_2 .

Let (A, B) be a separation of G such that $V(S_1) \subseteq A \setminus B$, $V(S_2) \subseteq B \setminus A$ and $X = A \cap B$. Let $f : V(G) \rightarrow \mathcal{L}$ be defined by

$$f(v) = \begin{cases} -1, & \text{if } v \in A \setminus B, \\ 0, & \text{if } v \in X, \\ 1, & \text{if } v \in B \setminus A. \end{cases}$$

Note that f is entire. By Lemma 5 we have

$$\int_{G \uparrow(i,j)} df = 2. \quad (1)$$

for $1 \leq i, j \leq n$.

For every $v \in V(G \uparrow(i,j)) \cap X$, let $\lambda(v) = \lambda_{G \uparrow(i,j),f}(v_i)$. For every $v \in X \setminus \bigcup_{1 \leq i,j \leq n} V(G \uparrow(i,j))$, let $\lambda(v) = 0$. By Lemma 8,

$$\int_{G \uparrow(i,j)} df = 2\lambda(V(G \uparrow(i,j)) \cap X) \quad (2)$$

It follows from (1) and (2) that $\lambda(X) = n^2$.

Let $H := G[X]$ and let $t := \text{tw}(H)$. We aim to prove that $t \geq \frac{n}{\sqrt{3\Delta}} - 1$. If $t > n^2/3 - 1$ or $t \geq n - 1$, then this claim holds, since $\Delta \geq 3$. Hence, we can assume that $n^2 \geq 3t + 3$ and $t < n - 1$. By Lemma 9 there exists a separation (K, L) of H such that

$$\frac{n^2}{3} \leq \lambda(K \setminus L) \leq \frac{2n^2}{3} \quad (3)$$

and $|K \cap L| \leq t + 1 < n$. Let $R := \{i \in [n] : V(R_i) \cap (K \cap L) = \emptyset\}$ and $C := \{j \in [n] : V(C_j) \cap (K \cap L) = \emptyset\}$, be the sets of indices of rows and, respectively, columns of (G, S_1, S_2) disjoint from $K \cap L$. Note that these sets are non-empty. Let $S := (R \times [n]) \cup ([n] \times C)$.

Let $g \in C^0(G, \mathcal{L})$ be defined by $g(v) = f(v)$ for $v \in V(G) \setminus L$ and $g(v) = \star$ for $v \in L$. Note that g is continuous and holomorphic on $V(G) - (K \cap L)$. Let $h(p) := \frac{1}{2} \int_{G \uparrow p} dg$ for $p \in [n]^2$. By Lemma 8, we have

$$h(p) = \lambda(V(G \uparrow p) \cap (K \setminus L)), \quad (4)$$

for every $p \in [n]^2$, and $h(p)$ is an integer for every $p \in [n]^2$ such that $V(G \uparrow p) \cap (K \cap L) = \emptyset$. Note that for $i_1 \in R$, $i_2 \in C$ and $1 \leq j_1, j_2 \leq n$, the paths $G \uparrow(i_1, j_1)$ and $G \uparrow(i_1, j_2)$ are g -homotopic, and the paths $G \uparrow(j_1, i_2)$ and $G \uparrow(j_2, i_2)$ are g -homotopic, since R_{i_1} and C_{i_2} are near-triangulations and g is holomorphic on them. It follows that $G \uparrow p_1$ and $G \uparrow p_2$ are g -homotopic for all $p_1, p_2 \in S$. Thus h is constant on S by Lemma 7. Let $H \in \mathbb{Z}$ be its value.

Note that $|[n] \setminus C| \leq t + 1$. Let us fix an element $c \in C$ and consider any $i \in [n]$. Let $t_i = |V(R_i) \cap (K \cap L)|$. For any $j \in [n] \setminus C$, the path $G \uparrow(i, j)$ is $(g, \Delta t_i)$ -almost homotopic to the path $G \uparrow(i, c)$, and by Lemma 7 we have $|h(i, j) - H| \leq \Delta t_i$. Thus

$$\left| \left(\sum_{j \in [n]} h(i, j) \right) - Hn \right| \leq \sum_{j \in [n]} |h(i, j) - H| \leq \Delta t_i(t+1). \quad (5)$$

Summing (5) over $i \in [n]$ and applying (4), we obtain

$$|\lambda(K \setminus L) - Hn^2| = \left| \sum_{p \in [n]^2} h(p) - Hn^2 \right| \leq \Delta(t+1)^2.$$

However, by (3), $\lambda(K \setminus L)$ differs from an integer multiple of n^2 by at least $n^2/3$. It follows that $t \geq \frac{n}{\sqrt{3\Delta}} - 1$, as desired. $\square$

3 Partitions of a cube

Consider the $N \times N \times N$ grid Q_N with non-decreasing diagonals. A path in Q_N with vertices $v_1 = (x_1, y_1, z_1), v_2 = (x_2, y_2, z_2), \dots, v_k = (x_k, y_k, z_k)$ in order is a *staircase* from v_1 to v_k if $x_i + 1 = x_{i+1}$, $y_i \leq y_{i+1}$ and $z_i \leq z_{i+1}$ for $1 \leq i \leq k-1$. The b -square $\square(v, b)$ around a vertex $v = (x, y, z) \in V(Q_N)$ is defined by

$$\square(v, B) = \{(x, y + d_y, z + d_z) : 0 \leq d_y, d_z \leq b\}.$$

The b -enlargement $B(b, P)$ of a staircase P is the subgraph of Q_N induced by $\bigcup_{v \in V(S)} \square(v, b)$. The *left side* of the b -enlargement of a staircase from u to v is $\square(u, b)$ and the *right side* is $\square(v, b)$. For each edge yz of the b -enlargement, let W_{yz} be the closed walk consisting of the path parallel to the staircase from the left side to y , the edge yz , the path parallel to the staircase from z to the left side, and possibly an edge in the left side.

We need the following property (another proof of a similar statement is implicit in [12], Proposition 3.1).

Lemma 10. *Let $b \geq 0$ be an integer, and let G be the b -enlargement of a staircase in Q_N . If $X \subseteq V(G)$ is a minimal set that intersects every path from the left side of G to the right side, then $G[X]$ is connected.*

Proof. Let S_1 and S_2 be the left and the right side of G , respectively. Since we can extend the staircase if necessary, we can assume that X is disjoint from $S_1 \cup S_2$. Let (A, B) be a separation of G such that $S_1 \subseteq A$, $S_2 \subseteq B$ and $X = A \cap B$. Let $l \in S_1$ and $r \in S_2$ be arbitrary vertices of the sides of G .

Consider any two vertices $x, y \in X$, and let C be the vertex set of the component of $G[X]$ containing x . Let $f : V(G) \rightarrow \mathcal{L}$ be defined by

$$f(v) = \begin{cases} -1, & \text{if } v \in A \setminus B, \\ 0, & \text{if } v \in C, \\ \star, & \text{if } v \in X \setminus C, \\ 1, & \text{if } v \in B \setminus A. \end{cases}$$

By the minimality of X , there exist paths P_x and P_y from l to r in G such that $V(P_x) \cap X = \{x\}$ and $V(P_y) \cap X = \{y\}$. Let W be the closed walk consisting of P_x and the reverse of P_y . Note that $I_W = \sum_{i=1}^n I_{T_i}$ for some triangles $T_1, \dots, T_n$ in G (since W is the sum of walks W_{yz} for edges yz of W together with a closed walk in the left side of G , it suffices to observe that this claim holds for the walks W_{yz} and for closed walks in the left side).

Since f is holomorphic, W is $(f, 0)$ -almost contractible, and thus P_x and P_y are f -homotopic. By Lemma 7, we have $\int_{P_x} df = \int_{P_y} df$. Since f is entire on P_x , we have $\int_{P_x} df = 2$, and thus $f(y) \neq \star$ (as otherwise we would have $\int_{P_y} df = 0$). Therefore, y is in the same component of $G[X]$ as x .

We conclude that $G[X]$ is connected. $\square$

Let A_1, A_2 be a partition of vertices of Q_N . Let $b \geq 0$ and $i \in \{1, 2\}$ be integers, and let P be a staircase. We say that P is (b, i) -blocked if every path in the b -enlargement of P joining its left side S_1 with its right side S_2 intersects $A_i \setminus (S_1 \cup S_2)$.

Lemma 11. *Let A_1, A_2 be a partition of vertices of Q_N . Let $b \geq 0$ and $i \in \{1, 2\}$ be integers. Let P be a (b, i) -blocked staircase, let M be the $(b+1)$ -enlargement of P , and let M_i be the subgraph of M induced by $A_i \cap V(M)$. There exists a connected component of M_i containing all paths in M_i which join the left side of M with the right side.*

Proof. Let M_0 be the b -enlargement of P . Let X be a minimal subset of $A_i \cap V(M_0)$ such that every path in M_0 from the left side to the right side intersects X . By Lemma 10, $M_0[X]$ is connected.

Let P' be a path in M_i joining the left side of M with the right side. For a vertex $u = (x, y, z) \in M$, let $v = (x, y_0, z_0) \in V(P)$ be the unique vertex of P such that $u \in \square(v, b+1)$. Let $\pi(u)$ be the point of $\square(v, b)$ closest to u in the Euclidean distance. That is, $\pi(u) = (x, \min(y, y_0 + b), \min(z, z_0 + b))$. Clearly u is adjacent to $\pi(u)$. Moreover, it is easy to check that if u and u' are adjacent then $\pi(u)$ and $\pi(u')$ are adjacent or equal.

It follows that the subgraph $\pi(P')$ of M_0 induced by $\{\pi(u) : u \in V(P')\}$ is connected and contains vertices both in the left and in the right side of M_0 . Therefore, $\pi(P')$ intersects X , and thus P' belongs to the same component of M_i as X , as desired. $\square$

The $t \times t$ -grid is the graph whose vertex set consists of all pairs $\{(x, y) \in \mathbb{Z}^2 : 0 \leq x, y \leq t-1\}$ and two vertices (x_1, y_1) and (x_2, y_2) are adjacent iff $|x_1 - x_2| + |y_1 - y_2| = 1$. Let P_1 and P_2 be two staircases. A staircase P joins P_1 with P_2 if P_1 is the initial segment of P and P_2 is the final segment of P , or vice versa.

Consider the $N \times N \times N$ grid Q_N . For an integer $n \geq 1$ and a point $v = (x, y, z) \in V(Q_N)$ with $x, y, z \leq N - n$, let $Q_n(v)$ denote the subgrid of Q_N induced by vertices $(x', y', z') \in V(Q_N)$ such that $x \leq x' \leq x + n - 1$, $y \leq y' \leq y + n - 1$, and $z \leq z' \leq z + n - 1$. For a positive integer d , let $p_d(j, k) = (4dj + 4dk, 2dj + dk, dj + 2dk)$. This definition is motivated by the following fact.

Observation 12. *Let $n \geq 1$, $b \geq 0$ and $d \geq n + b$ be integers. For every vertex $z = (j, k)$ of the $(2t + 1) \times (2t + 1)$ grid, let $p_z = p_d(j, k)$ and let P_z be a staircase in $Q_n(p_z)$. For every edge uz of the $(2t + 1) \times (2t + 1)$ grid, there exists a staircase P_{uz} joining P_u with P_z . Furthermore, the staircases can be chosen so that for any edges uz and $u'z'$ with $\{u, z\} \cap \{u', z'\} = \emptyset$, the b -enlargements of P_{uz} and $P_{u'z'}$ are vertex-disjoint.*

In the following proof of Theorem 1 we construct a subgraph of $Q_n[A_1]$ or $Q_n[A_2]$ which closely resembles a subdivision of the $t \times t$ grid. Unfortunately, obtaining an actual subdivision seem to require a lot more work, and therefore we content ourselves with obtaining the following less structured certificate of large tree-width. A collection of non-empty subsets $\mathcal{B}$ of the vertex set of a graph G is called a *bramble* if for all $B, B' \in \mathcal{B}$ the subgraph $G[B \cup B']$ of G induced by $B \cup B'$ is connected (and in particular, $G[B]$ is connected for every $B \in \mathcal{B}$). The *order* of $\mathcal{B}$ is the minimum size of the set $S \subseteq V(G)$ such that $S \cap B \neq \emptyset$ for every $B \in \mathcal{B}$. It is shown in [16] that if G contains a bramble of order t then G has tree-width at least $t - 1$.

We are now ready to complete the proof of Theorem 1.

Proof of Theorem 1. We will prove by induction on b that there exists $N = N(b)$ such that for every partition (A_1, A_2) of the vertex set of Q_N and each $i \in \{1, 2\}$, if the tree-width of both $Q_N[A_1]$ and $Q_N[A_2]$ is less than t , then Q_N contains a (b, i) -blocked staircase. Note that the theorem will follow, by choosing $b := \lceil \sqrt{18}(t + 1) \rceil - 1$ and $n := N(b)$. The b -enlargement B

of a staircase with sides S_1 and S_2 is a $(b+1, b+1)$ -slab. As every path in B from S_1 to S_2 intersects $A_i \setminus (S_1 \cup S_2)$, Theorem 4 implies that the tree-width of $Q_n[A_i]$ is at least $\frac{b+1}{\sqrt{18}} - 1 \geq t$.

The base case $b = 0$ is trivial with $N(0) = t + 2$: we have $V(Q_t(1, 1, 1)) \cap A_i \neq \emptyset$, since A_{3-i} induces a subgraph of tree-width less than t . We move on to the induction step. Let $n_0 = N(b-1)$ and let $N = (8t+5)(n_0+b)$. By the induction hypothesis, for every $v = (x, y, z)$ with $0 \leq x, y, z < N - n_0$, the subgrid $Q_{n_0}(v)$ contains a $(b-1, 3-i)$ -blocked staircase which we denote by P_v .

Let $d = n_0 + b$, and for every vertex $z = (j, k)$ of the $(2t+1) \times (2t+1)$ grid, let $p(j, k) = p_d(j, k)$. Note that $Q_{n_0}(p(j, k)) \subseteq Q_N$. Let $P_z = P(p(j, k))$; by Lemma 11, there exists a connected subgraph M_z of the b -enlargement B_z of P_z such that $V(M_z) \subseteq A_{3-i}$ and M_z contains all paths in $B_z[A_{3-i}]$ joining the left side of B_z with the right side.

For each edge yz of the $(2t+1) \times (2t+1)$ grid, let P_{yz} be the path as in Observation 12. If P_{yz} is (b, i) -blocked then the proof of the induction step is finished. Hence, we can assume that for every edge yz , there exists a path R_{yz} with $V(R_{yz}) \subseteq A_{3-i}$ in the b -enlargement of P_{yz} joining the left side of this enlargement to the right side. Note that R_{yz} intersects M_y and M_z .

For $0 \leq j \leq 2t$, let S'_j be the path forming the j -th column of the $(2t+1) \times (2t+1)$ grid, and let

$$S_j = V \left(\bigcup_{y \in V(S'_j)} M_y \cup \bigcup_{yz \in E(S'_j)} R_{yz} \right).$$

Let the set T_j be similarly defined for the j -th row of the grid. As observed in the previous paragraph, $Q_N[S_j]$ and $Q_N[T_j]$ are connected for all $0 \leq j \leq 2t$, and $M_{(j,k)} \subseteq S_j \cap T_k$. Let $\mathcal{B} = \{S_j \cup T_j\}_{0 \leq j \leq 2t}$. Clearly $\mathcal{B}$ is a bramble in $Q_N[A_{3-i}]$, and no vertex of Q_N belongs to more than two elements of $\mathcal{B}$ by Observation 12. It follows that the order of $\mathcal{B}$ is at least $t+1$, and thus $Q_N[A_{3-i}]$ has tree-width at least t , yielding the desired contradiction. $\square$

Acknowledgement. This research was partially completed at a workshop held at the Bellairs Research Institute in Barbados in April 2014. We thank the participants of the workshop, and especially Paul Seymour, for helpful discussions.

References

- [1] N. ALON, G. DING, B. OPOROWSKI, AND D. VERTIGAN, *Partitioning into graphs with only small components*, Journal of Combinatorial Theory, Series B, 87 (2003), pp. 231–243.
- [2] B. BAKER, *Approximation algorithms for NP-complete problems on planar graphs*, Journal of the ACM (JACM), 41 (1994), pp. 153–180.
- [3] C. CHEKURI AND J. CHUZHOUY, *Polynomial bounds for the grid-minor theorem*, in Proceedings of the 46th Annual ACM Symposium on Theory of Computing, STOC '14, New York, NY, USA, 2014, ACM, pp. 60–69.
- [4] J. CHUZHOUY, *Excluded grid theorem: Improved and simplified*, in Proceedings of the Forty-Seventh Annual ACM on Symposium on Theory of Computing, STOC '15, New York, NY, USA, 2015, ACM, pp. 645–654.
- [5] L. COWEN, W. GODDARD, AND C. E. JESURUM, *Defective coloring revisited*, Journal of Graph Theory, 24 (1997), pp. 205–219.
- [6] M. DEVOS, G. DING, B. OPOROWSKI, D. SANDERS, B. REED, P. SEYMOUR, AND D. VERTIGAN, *Excluding any graph as a minor allows a low tree-width 2-coloring*, J. Comb. Theory, Ser. B, 91 (2004), pp. 25–41.
- [7] G. DING, B. OPOROWSKI, D. P. SANDERS, AND D. VERTIGAN, *Surfaces, tree-width, clique-minors, and partitions*, Journal of Combinatorial Theory, Series B, 79 (2000), pp. 221–246.
- [8] Z. DVOŘÁK, *Sublinear separators, fragility and subexponential expansion*, European Journal of Combinatorics, 52, Part A (2016), pp. 103–119.
- [9] K. EDWARDS, D. Y. KANG, J. KIM, S.-I. OUM, AND P. SEYMOUR, *A relative of hadwiger’s conjecture*, ArXiv e-prints, 1407.5236 (2014).
- [10] D. EPPSTEIN, *Diameter and treewidth in minor-closed graph families*, Algorithmica, 27 (2000), pp. 275–291.
- [11] C. LIU AND S. OUM, *Partitioning h -minor free graphs into three subgraphs with no large components*, Electronic Notes in Discrete Mathematics, 49 (2015), pp. 133–138.

- [12] J. MATOUŠEK AND A. PŘÍVĚTIVÝ, *Large monochromatic components in two-colored grids*, SIAM Journal on Discrete Mathematics, 22 (2008), pp. 295–311.
- [13] N. ROBERTSON AND P. D. SEYMOUR, *Graph Minors. III. Planar tree-width*, J. Combin. Theory, Ser. B, 36 (1984), pp. 49–64.
- [14] —, *Graph Minors. V. Excluding a planar graph*, J. Combin. Theory, Ser. B, 41 (1986), pp. 92–114.
- [15] N. ROBERTSON, P. D. SEYMOUR, AND R. THOMAS, *Quickly excluding a planar graph*, J. Combin. Theory, Ser. B, 62 (1994), pp. 323–348.
- [16] P. D. SEYMOUR AND R. THOMAS, *Graph searching and a min-max theorem for tree-width*, Journal of Combinatorial Theory, Series B, 58 (1993), pp. 22–33.